# Developing Objective Drought Indicator Blends Using Principal Component Analysis

Tsegaye Tadesse, Ph.D.
Research Associate Professor,
Climatologist and Remote Sensing Expert,
Geospatial Coordinator
National Drought Mitigation Center (NDMC)
University of Nebraska-Lincoln, USA



## Acknowledgement

## This work is supported by NDMC's USDA- and NOAA-funded projects Research Team:

Tsegaye Tadesse, Geospatial Coordinator, NDMC

Mark Svoboda, Director, NDMC

Brian Fuchs, Monitoring Coordinator, NDMC, UNL

Yared Ashenafi, Research Scientist, NDMC, UNL

Getachew Berhan, Post-doctoral Researcher, NDMC, UNL


Chris Poulsen, GIS Manager, NDMC, UNL

John Swigart, Geospatial Analyst, NDMC, UNL

Jeff Nothwehr, GIS and Web Specialist, NDMC, UNL

Brian Wardlow, Director, CALMIT, UNL

Andualem Shiferaw, Graduate Research Assistant, NDMC, UNL





We would like to have collaborators from the drought community who are interested!!

## Outline

- 1. Introduction: Objective Drought Indicator blend (ODI-blend)
  - Current method (NOAA-CPC)
  - PCA-based weighting
- 2. Current short- and long-term ODI blends
  - Data Inputs and historical maps with higher spatial resolution
- 3. Developing short- and long-term ODI blends using PCA
  - Data Inputs, Modeling, Evaluation, and Research Results
- 4. Summary and the way forward

## Methods to Develop Objective Drought Indicators (ODI) Blends

**Goal:** Objectively combining/blending drought indices based on the potential contribution of each input variables to drought.

### Methods of components weighting


- 1. Domain Expert judgment (NOAA-CPC's Experimental Objective Blends of Drought Indicators)
- 2. Weights based on statistical models: Principal Component Analysis (PCA)

#### PCA-based USDM objective blend maps (PCA-based USDM)

 The main objective of this research is to provide an alternative objective weighting scheme (PCA approach) for the USDM blend maps and improve its spatial resolution.

### **Current short- and long-term ODI blends**

Weights are based on experts judgments




<sup>\*</sup>These weights are drought experts' judgment inputs and weightings for the short- and long-term objective blend maps.

<sup>\*\*</sup>NOAA used CPC soil moisture model. Because of data availability, we used NLDAS-Noah Soil moisture data.

### Historical short- and long-term ODI blends

(based on expert judgments short-term objective blend)

\*\* NDMC produced historical ODI blend maps for the CONUS (1980-2012) at a higher resolution (~12.5 km)



#### Expert-determined input components and weightings\*:


We have produced 3,432 historical (1980-2012) ODI blend maps similar to the NOAA-CPC climate-division blends, but at 12.5 km resolution.

<sup>\*</sup>These weights are drought experts' judgment inputs and weightings for the short- and long-term objective blend maps.


<sup>\*\*</sup>NLDAS-Noah Soil moisture data.

<sup>\*\*\*</sup> Input maps in the fig. above are for illustration purpose only.


# PCA-based Combined Drought Indices (CDIs) Methodological Approach




PCA-based Percentage contributions of each input parameter for the short-term USDM blend




### 2011 Short-term Drought Indicator Blend




### 2012 Short-term Drought Indicator Blend




Week 39:Sept23






PCA-based Percentage contributions of each input parameter for the long-term Eastern\* US blend **PHDI** SPI-24 Week 21/May 21 SM SPI-12 SPI-6 **SPI-60** *Areal avg.* = 27% Areal avg. = 23% **Areal avg. = 21%** Areal avg. = 14% *Areal avg. = 27%* Areal avg. = 13% Week 31/July 28 *Areal avg. = 26%* Areal avg. = 22% Areal avg. = 23% Areal avg. = 26% **Areal avg. = 15%** Areal avg. = 12% Week 39/Sept23 *Areal avg. = 26%* Areal avg. = 22% **Areal avg. = 26%** Areal avg. = 23% Areal avg. = 14% **Areal avg. = 12%** 20 - 30 30 - 40 10 - 20 0 - 10 50 - 60 40 - 50 Eastern\* US blend Long-term objective blend maps Experts' weight 60-month SPI 24-month SPI 12-month SPI 6-month SPI SM-Noah\*\* **PHDI** (10%)(20%)(20%)(15%)(10%)(25%)

### 2011 Long-term Drought Indicator Blend




### 2012 Long-term Drought Indicator Blend



### Future works: blending more input data

New input historical datasets for new blends maps may include:


- QuickDRI
- VegDRI
- ESI
- EDDI
- VPD
- GRACE



- ■Short and long-term blends using:
  - longer time series data (1980-2017)
  - shorter time series (2003-2017)

### **USDM** Blends at the NDMC website

(in development: http://dmcommunity.unl.edu/USDMBlends.aspx)



\*Requires user name and password

### Summary

☐ We've recreated blend maps and compared with the NOAA CPC's experimental blend maps (sanity check) Assessed the blends using selected years (e.g., 2011 and 2012) in the growing season Reasonably similar drought patterns and intensity observed Developed short- and long-term ODI blends using PCA weighting scheme Avoids/limits the subjectivity of experts' judgment (though it's educated guess) ☐ Improved the spatial resolution New PCA-based blends preserve the seasonality of the climatic and/or growing periods ☐ Each week in a year has a unique weights for the inputs as opposed to a "one-fits-all" weighting scheme We've produced historical short- and long-term ODI maps for CONUS (using both weighting schemes) □ Short-term weekly ODI (1980-2012) – 3432 maps □Long-term weekly ODI (1980-2012) – 3432 maps ☐ All blend maps do have higher resolution (i.e., 12.5 km vs climate division NOAA-CPC blends) ☐ Are available at the NDMC website (but not publicly available yet) More evaluation with ground observation is needed We're exploring to include new input satellite-derived data and hybrid drought indices.

## Thank You



#### Tsegaye Tadesse, Ph.D.

Research Associate Professor, Climatologist and Remote Sensing Expert Geospatial Science Coordinator National Drought Mitigation Center (NDMC) University of Nebraska-Lincoln, USA

Email: ttadesse2@unl.edu

DROUGHT.UNL.EDU

e | ndmc@unl.edu





